

POLYDIFFERENTIALS AND THE DEFORMATION FUNCTOR OF CURVES WITH AUTOMORPHISMS II

A. KONTOGEORGIS

ABSTRACT. We apply the known results on the Galois module structure of the sheaf of polydifferentials in order to study the dimension of the tangent space of the deformation functor of curves with automorphisms. We are able to find the dimension for the case of weakly ramified covers and for the case of the action of a cyclic group of order p^v .

1. INTRODUCTION

Let X be a non-singular complete curve of genus $g \geq 2$ defined over an algebraic closed field k of positive characteristic $p > 3$, and let G be a subgroup of the automorphism group of X . In [1] J.Bertin, A. Mézard proved that the equivariant cohomology of Grothendieck $H^1(G, \mathcal{T}_X)$ is the tangent space of the global deformation functor of smooth curves with automorphisms. The dimension of the k -vector space $H^1(G, \mathcal{T}_X)$ is a measure of the directions a curve can be deformed together with a subgroup of the automorphism group. The computation of this dimension turns out to be a difficult problem, and in the literature there are results concerning ordinary curves [5], cyclic groups of order p [1]. In [9] the author attempts to compute this space using the low terms of the Lyndon-Hochschild-Serre spectral sequence. This allows us to compute the dimension of $H^1(G, \mathcal{T}_X)$ if G is an elementary abelian group *i.e.* isomorphic to a direct product of cyclic groups of order p . The disadvantage of this method is that it involves the computation of the transfer map that depends on the group structure of the decomposition series of the decomposition groups $G(P)$ at the wild ramification points.

The author in [10] proposed an alternative method of approaching the problem of computing the dimension of $H^1(G, \mathcal{T}_X)$. Precisely one can prove that for a p -group G and for the sheaf Ω_X of differentials on X the following equation holds:

$$H^1(G, \mathcal{T}_X) = H^0(X, \Omega_X^{\otimes 2})_G.$$

This observation allows us to compute the desired dimension if we know the Galois module structure of the space of 2-holomorphic differentials. The knowledge of the Galois module structure of the space of holomorphic differentials in positive characteristic is still an open problem and there are only partial results in the case of tame covers [7],[15],[12] and when G is a cyclic group of order equal to the characteristic [14]. The computation of the Galois module structure of holomorphic differentials in the tame case does not offer any new information for the computation of $H^1(G, \mathcal{T}_X)$ since the classical methods from the theory of Riemann surfaces can be used in this case.

In [10] the author was able to recover the result of J.Bertin, A. Mézard [1] concerning the dimension of $H^1(G, \mathcal{T}_X)$ in the case $G = \mathbb{Z}/p\mathbb{Z}$ by applying the computation of Nakajima [14] concerning the $\mathbb{Z}/p\mathbb{Z}$ -module structure of $\Omega_X^{\otimes 2}$.

In this article we use the results of B. Köck [8] on weakly ramified covers, *i.e.* on covers so that $G_2(P) = \{1\}$ for all wild ramified points, in order to determine $\dim_k H^1(G, \mathcal{T}_X)$. Since all ordinary curves are weakly ramified [16, th. 2i] we recover the result of Cornelissen and Kato [5] for deformations of ordinary curves.

Many authors [25],[13],[7],[3],[24], in order to study the $k[G]$ -module structure of spaces $\Omega_X(-D)$ considered the $k[G]$ -module structures of the spaces of semisimple and nilpotent differentials with respect to the Cartier operator. In section 4 we follow this approach. The representation module corresponding to semisimple differentials is studied and the problem of computing the space of covariant differentials is reduced to the problem of computing nilpotent covariant differentials. More precisely we are able to prove that:

$$(1) \quad \dim_k H^1(G, \mathcal{T}_X) = 2(g_Y - 1) + \gamma_Y - 1 + r + \dim_k \Omega_X(-D)_G^n,$$

where g_Y, γ_Y denote the genus and the p -rank of the Jacobian of $Y = X/G$ and r is the number of points of Y ramified in $X \rightarrow X/G$.

In his PhD thesis [2] N. Borne proved an equivariant Riemann-Roch theorem and provided an equivariant Euler characteristic $K_0(G, X) \rightarrow R_k(G)$, where $K_0(G, X)$ is the Grothendieck group of $G - \mathcal{O}_X$ -modules and $R_k(G)$ is the Brauer Character Group. In the case of p -groups in positive characteristic this approach does not give us more information than the classical Riemann-Roch theorem; the group $R_k(G) = \mathbb{Z}$ [21, 15.6] and equivariant information is lost.

N. Borne in [4] proposed a refinement of his equivariant K -theory by using the ideas of Auslander in modular representation theory. This new theory enables him to compute the Galois module structure of spaces of global sections of linear systems of curves of the form $\mathcal{O}_X(D)$ where $\deg D \geq 2g_X - 2$ and G is a cyclic group of the form $\mathbb{Z}/p^v\mathbb{Z}$. We apply the results of Borne in order to prove:

Proposition 1.1. *Assume that the group $G = \mathbb{Z}/p^v\mathbb{Z}$ acts on X . Let X_{ram} denote the set of ramification points of the cover $\pi : X \rightarrow X/G = Y$. For every point $P \in X_{ram}$ let N_P denote the highest jump in the lower ramification filtration and $f(P)$ the highest jump in the upper ramification filtration. Let $\Delta : \mathbb{Z} \rightarrow \mathbb{Z}$ denote the map $\Delta : a \mapsto [a/p]$, and Δ^k the composition of k times the map Δ . Let g_Y be the genus of the quotient curve Y and let $r = \#\pi_*(X_{ram})$.*

The dimension of $H^1(\mathbb{Z}/p^v\mathbb{Z}, \mathcal{T}_X)$ is given by:

$$\dim_k H^1(\mathbb{Z}/p^v\mathbb{Z}, \mathcal{T}_X) = 3(g_Y - 1) + 2r + \sum_{P \in \pi_*(X_{ram})} (2f(P) + \Delta^k(-2N_P - 2)).$$

This result can be obtained also by using Grothendieck's equivariant theory. One can compute the the group $H^0(Y, \pi_*^G(\mathcal{T}_X))$ [1],[9, sec 3.] and the local contributions at each wild ramified point given in [1, prop. 4.1.1].

In deformation theory it is an interesting problem to compare the dimension of the tangent space of the deformation functor to the dimension of the versal deformation ring. In the case of ordinary curves and in the p -cyclic case the Krull dimension has been computed [5],[1]. In [17] R. Pries studied unobstructed deformations of curves acted on by Abelian groups under the additional assumption that the deformations do not split the branch locus. A comparison of the result of

Pries to the result of proposition 1.1 gives the dimension of the space generated by obstructed deformations and of deformations that split the branch locus.

Acknowledgements The author would like to thank G. Cornelissen and M. Matignon for pointing him to the work of N. Borne, B. Köck, N. Stalder. The author would like to thank B. Köck for his remarks and corrections on an earlier version of this article.

2. SPACES OF 2-HOLOMORPHIC DIFFERENTIALS.

We will denote by Ω_X the sheaf of differentials of X . Let D be an effective divisor on the curve X and let G be a finite subgroup of $\text{Aut}(X)$. The set of meromorphic differentials on X is denoted by $\mathcal{M}_X = \{xdy : x, y \in K(X)\}$, where $K(X)$ denotes the function field of the curve X . We will denote by

$$\Omega_X(D) = \{\omega \in \mathcal{M}_X : \text{div}(\omega) \geq D\} = H^0(X, \Omega_X \otimes \mathcal{O}_X(-D)).$$

and by

$$\mathcal{L}_X(D) = \{f \in K(X) : \text{div}(f) \geq -D\} = H^0(X, \mathcal{O}_X(D)).$$

We will denote the k -dimension of $\mathcal{L}_X(D)$ by $\ell(D)$. The reader should be careful about the notation here. We follow the notation of Serre, used also in the papers of Stichtenoth, Nakajima, Subrao. In the papers of N. Borne [4], E. Köck [8], E. Kani [7], N. Stalder [24] the notation $\Omega_X(D)$ is used for what we write as $\Omega_X(-D)$.

Lemma 2.1. *Let G be a p -group and consider the cover $X \rightarrow X/G$. We assume that the genus of the curve X is $g_X \geq 2$. There is a G -invariant differential ω in X , such that $\text{div}(\omega)$ is effective.*

Proof. Let $b_1, \dots, b_r$ be the ramification points of the cover $\pi : X \rightarrow Y = X/G$. In order to find ω we have to select a meromorphic differential ϕ of Y and take the pullback $\pi^*(\phi)$. Then $\pi^*(\phi)$ is G -invariant and we have to select it so that $\text{div}(\phi) \geq 0$. Fix a meromorphic differential ϕ_1 on Y and consider the set $f\phi_1$ where f is an arbitrary element in the function field of Y . We have that

$$(2) \quad \text{div}(\pi^*(f\phi_1)) = \pi^*\text{div}(f\phi_1) + R \geq 0,$$

where R is the ramification divisor given by

$$R = \sum_{i=1}^r \sum_{P \mapsto b_i} \sum_{\nu=0}^{\infty} (e_{\nu}(P) - 1)P.$$

In order to check that the divisor on the left hand side of (2) is positive we push forward again and arrive at

$$(3) \quad \text{div}(f\phi_1) + \sum_{i=1}^r \sum_{\nu=0}^{\infty} \frac{e_{\nu}(b_i) - 1}{e_0(b_i)} \geq 0.$$

The later condition is equivalent to $f \in \mathcal{L}_Y(K + A)$, where A is the effective divisor

$$A := \sum_{i=1}^r \left\lfloor \sum_{\nu=0}^{\infty} \frac{e_{\nu}(b_i) - 1}{e_0(b_i)} \right\rfloor b_i.$$

Using Riemann-Roch on the curve Y we compute

$$\ell(K + A) = \ell(-A) + 2g_Y - 2 + \deg(A) - g_Y + 1 = g_Y - 1 + \deg(A) \geq 1.$$

Indeed, since G is a p -group and $g_X > 2$ there should be at least one wild ramified point. Every ramified point contributes at least 1 to the degree of A , since $e_0(b_i) =$

$e_1(b_i)$ and if $g_Y \geq 1$ we are done. In the case $g_Y = 0$ we use $g_X \geq 2$ and the Riemann-Hurwitz formula to see that there should be either at least two ramified points so $\deg A \geq 2$, or one ramified point with $\sum_{\nu=0}^{\infty} \frac{e_{\nu}(b_i)-1}{e_0(b_i)} > 2$, and in this case also $\deg A \geq 2$.

Every $f \in \mathcal{L}_Y(K+A)$ gives rise to the desired differential $\omega = \pi^*(f\phi_1)$. Moreover we can select $f \in \mathcal{L}_Y(K+A)$ such that it has polar divisor A . This implies that the support of $\pi^*(f\phi_1)$ has no intersection with the branch locus. $\square$

Proposition 2.2. *Assume that the p -group G acts on the group G . There is an effective and invariant divisor $D^* = \operatorname{div}(\pi^*(\phi)) = \pi^*(\operatorname{div}(\phi)) + R \geq 0$ so that the module $\Omega_X(-D^*)$ is isomorphic to $H^0(X, \Omega_X^{\otimes 2})$ as a $k[G]$ -module.*

Proof. According to lemma 2.1 we can select ϕ so that $\pi^*(\phi)$ is G -invariant and the divisor $D^* := \operatorname{div}(\pi^*(\phi)) = \pi^*(\operatorname{div}(\phi)) + R$ is an effective G -invariant divisor. Every differential can be written as $f\phi$ and every 2-differential is an expression of the form $\omega \otimes \phi$, for an other differential ω on X . The space

$$\begin{aligned} H^0(X, \Omega_X^{\otimes 2}) &= \{\omega \otimes \phi : \operatorname{div}(\omega \otimes \phi) \geq 0\} = \\ &= \{\omega \in \Omega_X : \operatorname{div}(\omega) \geq -\operatorname{div}(\phi)\}. = \\ &= \Omega_X(-D^*). \end{aligned}$$

$\square$

3. THE WEAKLY RAMIFIED CASE

In this section we assume that the cover $X \rightarrow X/G$ is *weakly* ramified, i.e. for every $P \in S$ we have that $G_i(P) = \{1\}$ for all $i \geq 2$. The group G is always assumed to be a p -group. The ramification divisor is computed $R = \sum_{P \in S} 2(e_0(P) - 1)P$.

Lemma 3.1. *The module $\Omega_X(-D^* - \sum_{P \in S} 3P)$ is a projective $k[G]$ -module.*

Proof. Let K be a canonical divisor. Let us write

$$\Omega_X(-D^* - \sum_{P \in S} 3P) = \mathcal{L}_X(K + D + \sum_{P \in S} 3P).$$

Let $D' := K + D^* + \sum_{P \in S} 3P = \sum n_P P$. We compute that

$$D' = K + D^* + \sum_{P \in S} 3P = 2\pi^*(\operatorname{div}(\phi)) + \sum_{P \in S} 4(e_0(P) - 1) + 3P.$$

Since we have assumed that G is a p -group we have that $e_0(P) = e_1(P) = e^w(P)$. Therefore $n_P \equiv -1 \pmod{p}$ for all wild ramified points of $X \rightarrow X/G$. On the other hand

$$H^1(X, \mathcal{O}_X(D')) = \mathcal{L}_X(K - D') = 0,$$

since $\deg(K - D') < 0$. The desired result follows by [8, th. 2.1].

B. Köck proposed to me the following more abstract approach: By [2, th. 44] there is a G -invariant canonical divisor $K_X = \sum_P m_P P$ on X . The proof of corollary 2.3 in [8] implies that all $m_P \equiv -2 \pmod{p}$. Therefore the divisor, $2K_X + \sum_{P \in S} 3P = \sum n_P P$ has $n_P \equiv 2(-2) + 3 = -1 \pmod{p}$ as required. $\square$

We can now form the short exact sequence:

$$(4) \quad 0 \rightarrow \Omega_X(-D^*) \rightarrow \Omega_X(-D^* - \sum_{P \in S} 3P) \rightarrow \frac{\Omega_X(-D^* - \sum_{P \in S} 3P)}{\Omega_X(-D^*)} \rightarrow 0.$$

The short exact sequence of sheaves

$$0 \rightarrow \Omega_X \otimes \mathcal{O}_X(-D^*) \rightarrow \Omega_X \otimes \mathcal{O}_X(-D^* - \sum_{P \in S} 3P) \rightarrow \Sigma \rightarrow 0$$

where

$$\Sigma := \frac{\Omega_X \otimes \mathcal{O}_X(-D^* - \sum_{P \in S} 3P)}{\Omega_X \otimes \mathcal{O}_X(-D^*)}$$

gives rise to a long exact sequence of k -vector spaces by applying the functor of global sections. This long exact sequence combined with $H^1(X, \Omega_X \otimes \mathcal{O}_X(-D^*)) = 0$ allows us to express

$$H^0(X, \Sigma) = \frac{\Omega_X(-D^* - \sum_{P \in S} 3P)}{\Omega_X(-D^*)}$$

as the direct sum of the stalks Σ_P of the skyscraper sheaf Σ at points $P \in S$. Let us denote by $\Sigma' := H^0(X, \Sigma)$.

Equation (4) gives the following long exact sequence:

$$0 \rightarrow H_1(G, \Sigma') \rightarrow \Omega_X(-D^*)_G \rightarrow \Omega_X(-D^* - \sum_{P \in S} 3P) \rightarrow \Sigma'_G \rightarrow 0,$$

since $\Omega_X(-D^* - \sum_{P \in S} 3P)$ is a $k[G]$ -projective module. Therefore, the desired dimension can be computed:

$$(5) \quad \dim_k \Omega_X(-D^*)_G = \dim_k H_1(G, \Sigma') + \dim_k \Omega_X(-D^* - \sum_{P \in S} 3P)_G - \dim_k \Sigma'_G$$

In what follows we will compute every summand on the right hand side of (5).

Lemma 3.2. $\dim_k \Omega_X(-D^* - \sum_{P \in S} 3P)_G = 3(g_Y - 1) + 3r$.

Proof. Using the theorem of Riemann-Roch we compute:

$$(6) \quad \dim_k \Omega_X(-D^* - \sum_{P \in S} 3P) = 2g_X - 2 + |G|(2g_Y - 2) + \sum_{P \in S} (2(e_0(P) - 1) + 3) \\ - g_X + 1 + \dim_k \mathcal{L}_X(-D^* - \sum_{P \in S} 3P).$$

But $\deg(-D^* - \sum_{P \in S} 3P) < 0$ therefore $\dim_k \mathcal{L}_X(-D^* - \sum_{P \in S} 3P) = 0$. Riemann-Hurwitz implies that

$$(7) \quad 2g_X - 2 = |G|(2g_Y - 2) + 2 \sum_{P \in S} (e_0(P) - 1).$$

By combining (6),(7) we obtain:

$$\dim_k \Omega_X(-D^* - \sum_{P \in S} 3P) = 3|G|(g_Y - 1) + 3r|G|.$$

Since $\Omega_X(-D^* - \sum_{P \in S} 3P)$ is projective it is of the form $k[G]^a$, where $a = 3(g_Y - 1) + 3r$, and each $k[G]$ direct summand contributes 1 to the dimension of $\Omega_X(-D^* - \sum_{P \in S} 3P)_G$. The desired result follows. $\square$

Let us now study the space Σ' as a G -module: The differential $\pi^*(\phi)$ defined in lemma 2.1 is an invariant differential and we have that $\text{div}(\pi^*(\phi)) = D^*$. Every differential ω can be written as $\omega = f\pi^*(\phi)$, Therefore,

$$\Omega_X(-D^*) = \{\text{div } f\pi^*(\phi) > -D^*\} \cong \mathcal{O}_X(2D^*),$$

and

$$\Omega_X(-D^* - \sum_{P \in S} 3P) \cong \mathcal{O}_X(2D^* + \sum_{P \in S} 3P),$$

where the last two isomorphisms are isomorphisms of $k[G]$ -modules.

Let $O(P) = \{gP : g \in G\}$ denote the orbit of P under the action of the group G . The set S of ramification points can be written as a disjoint union of orbits of points of X :

$$S = \bigcup_{j=1}^r O(P_j),$$

for a selection $P_1, \dots, P_r$ of points of X . We can write $H^0(X, \Sigma)$ as

$$(8) \quad \Sigma' = \bigoplus_{j=1}^r \bigoplus_{P \in O(P_j)} \Sigma_P,$$

Σ_P is the stalk of Σ at P . Thus, (8) can be written as

$$\Sigma' = \bigoplus_{j=1}^r \text{Ind}_{G(P_j)}^G \Sigma_{P_j}.$$

Shapiro lemma [26, 6.3.2, p.171] implies that

$$(9) \quad H_*(G, \Sigma') = \bigoplus_{j=1}^r H_*(G, \text{Ind}_{G(P_j)}^G \Sigma_{P_j}) = \bigoplus_{j=1}^r H_*(G(P_j), \Sigma_{P_j}).$$

Let P be a ramified point. We have assumed that the cover is weakly ramified we have that $G_2(P) = \{1\}$. This implies that for a local uniformizer t_P at P we have

$$(10) \quad G(P) \ni g : t_P \mapsto t_P \left(1 + \alpha_1(g)t_P + \sum_{\nu=2}^{\infty} \alpha_{\nu}(g)t_P^{\nu} \right),$$

where $\alpha_1(g) \neq 0$. The map $g \mapsto a_1(g)$ is a homomorphism and allows us to see $G(P)$ as a finite dimensional $\mathbb{F}_p$ -vector subspace of k .

The quotient Σ_P is then generated as a k -vector space by elements

$$\Sigma_P = \left\{ \omega_1 := \frac{1}{t_P^{4e(P)-1}}, \omega_2 := \frac{1}{t_P^{4e(P)-2}}, \omega_3 := \frac{1}{t_P^{4e(P)-3}} \right\}$$

and the action is given by:

$$(11) \quad \sigma : \frac{1}{t_P^{4e(P)-1}} \mapsto \frac{1 + \alpha_1(g)t_P + \alpha_2(g)t_P^2}{t_P^{4e(P)-1}} = \frac{1}{t_P^{4e(P)-1}} + \alpha_1(g) \frac{1}{t_P^{4e(P)-2}} + \alpha_2(g) \frac{1}{t_P^{4e(P)-3}},$$

since $\binom{-4e(P)+1}{1} = 1$ and $\binom{-4e(P)+1}{2} = 0$.

$$(12) \quad \sigma : \frac{1}{t_P^{4e(P)-2}} \mapsto \frac{1 + 2\alpha_1(g)t_P}{t_P^{4e(P)-2}} = \frac{1}{t_P^{4e(P)-2}} + 2\alpha_1(g) \frac{1}{t_P^{4e(P)-3}}$$

and

$$(13) \quad \sigma : \frac{1}{t_P^{4e(P)-3}} \mapsto \frac{1}{t_P^{4e(P)-3}}.$$

Lemma 3.3. *We have that $\dim_k \Sigma'_G = H_0(G, \Sigma') = r$*

Proof. Let P be one of the $\{P_1, \dots, P_r\}$. Let $\omega \in \Sigma_P$. Using equations (11),(12),(13) we observe that there is only one invariant element in Σ_P , therefore the images of the linear maps $(g-1)\omega$ are all isomorphic and two dimensional generated by the elements ω_2, ω_3 and so we arrive at:

$$\dim_k H_0(G(P), \Sigma_P) = \dim_k \frac{\Sigma_P}{g\omega - \omega} = 1.$$

For the global computation we use (9) in order to obtain:

$$\dim_k \Sigma'_G = H_0(G, \Sigma') = r.$$

□

Lemma 3.4. *We have that $\dim_k H_1(G, \Sigma') = \sum_{j=1}^r \log_p |G(P_j)| - r$*

Proof. Let P be one of the $\{P_1, \dots, P_r\}$. We will use the *normalised bar resolution* defined in [26, 6.5, p. 177] in order to compute $H_1(G, \Sigma_P)$. Recall that $k[G(P)] \cong B_0$ and B_0 is generated by the symbol $[\]$, B_1 is the free $k[G(P)]$ -module on the set of symbols $\{[g] : g \in G(P) \setminus \{1\}\}$, and B_2 is the free $k[G(P)]$ -module on the set of symbols $\{[g|h] : g, h \in G(P) \setminus \{1\}\}$. For the differential maps we have $\partial_1 : B_1 \rightarrow B_0$ and $\partial_1[g] = (g-1)[\]$, and $\partial_2 : B_2 \rightarrow B_1$, $\partial_2[g|h] = g[h] - [gh] + [g]$. Higher B_n can be similarly defined but we don't need them here. The group $H_1(G, \Sigma_P)$ is defined by the homology at position 1 of the chain complex $\Sigma_P \otimes B_*$. We have:

$$(14) \quad H_1(G, \Sigma_P) = \frac{\ker(\partial_1 : \Sigma_P \otimes B_1 \rightarrow \Sigma_P \otimes B_0)}{\text{im}(\partial_2 : \Sigma_P \otimes B_2 \rightarrow \Sigma_P \otimes B_1)}.$$

We will focus first on the study of the local components Σ_P . Let $\omega_3 = \frac{1}{t_P^{4e(P)-3}}$, $\omega_2 = \frac{1}{t_P^{4e(P)-2}}$, $e_1 = \frac{1}{t_P^{4e(P)-1}}$. Observe that the space generated by $\omega_3 \otimes [g]$ is in $\ker \partial_1$, since

$$\partial_1(\omega_3 \otimes [g]) = \omega_3(g-1) \otimes [\] = 0.$$

On the other hand

$$\begin{aligned} \partial_2(\omega_2 \otimes [g|h]) &= \omega_2 \otimes (g[h] - [gh] + [g]) = \omega_2 g \otimes [h] - \omega_2 \otimes [gh] + \omega_2 \otimes [g] = \\ &= \alpha_1(g) \omega_3 \otimes [h] + \omega_2 \otimes (g[h] - [gh] + [g]). \end{aligned}$$

Since ω_2, ω_3 are linear independent and $\alpha_1(g) \neq 0$, we obtain that all elements of the form $\omega_3 \otimes [h]$, $h \in G(P)$ are 0 in the homology group $H_1(G(P), \Sigma_P)$.

Observe that an element $\sum_{g \in G(P)} \lambda_g \omega_2 \otimes [g]$ is in the kernel of ∂_1 if and only if $\sum_{g \in G(P)} \lambda_g \alpha_1(g) = 0$, since we have assumed that $p \neq 2$. On the other hand

$$(15) \quad \partial_2(\omega_1 \otimes [g|h]) = \omega_1 \otimes (g[h] - [gh] + [g]) = \omega_2 \alpha_1(g) \otimes [h] + \omega_1([h] - [gh] + [g]),$$

therefore $\omega_2 \otimes [h]$ is zero in the homology group $H_1(G, \Sigma_P)$.

Consider an element

$$\omega = \sum_{g \in G(P)} \lambda_g \omega_1 \otimes [g] + \sum_{g \in G(P)} \mu_g \omega_2 \otimes [g] + \sum_{g \in G(P)} \nu_g \omega_3 \otimes [g],$$

and the image $\partial_1 \omega$ given by

$$(16) \quad \partial_1(\omega) = \sum_{g \in G(P)} \alpha_1(g) \lambda_g \omega_2 \otimes [\] + \sum_{g \in G(P)} (2\alpha_2(g) \lambda_g + \mu_g \alpha_1(g)) \omega_3 \otimes [\].$$

Equation (16) gives two necessary conditions for $\omega \in \ker \partial_1$, namely:

$$\sum_{g \in G(P)} (2\alpha_2(g)\lambda_g + \mu_g\alpha_1(g)) = 0$$

and

$$\sum_{g \in G(P)} \alpha_1(g)\lambda_g = 0.$$

We have seen that $H_1(G(P), \Sigma_P)$ is generated by the images of elements of the form $\omega_1 \otimes [g]$. On the other hand, equation (15) implies that:

$$(17) \quad \omega_1 \otimes [gh] = \omega_1 \otimes [g] + \omega_1 \otimes [h] \text{ in } H_1(G(P), \Sigma_P).$$

The groups $G(P)$ are elementary Abelian and can be written as $G(P) = \bigoplus_{i=1}^t g_i \mathbb{F}_p$. Equation (17) implies that

$$H_1(G, \Sigma_P) \subset \omega_1 \otimes \langle [g_i] : i = 1, \dots, t \rangle_k.$$

For a linear combination $\sum_{i=1}^t \lambda_i \omega_1 \otimes [g_i]$ we have that it is in $\ker \partial_1$ if and only if $\sum_{i=1}^t \lambda_i = 0$. Therefore:

$$\dim_k H_1(G, \Sigma_P) = \log_p |G(P)| - 1.$$

The global contribution is computed using (9) and equals to

$$H_1(G, \Sigma') = \sum_{j=1}^r \log_p |G(P_j)| - r,$$

i.e., the desired result. $\square$

Observe that the dimension of the tangent space of the deformation functor can now be computed by combining (5) and lemmata 3.2, 3.3, 3.4:

$$\dim_k H^1(G, \mathcal{T}_X) = 3g_Y - 3 + r + \sum_{j=1}^r \log_p |G(P_j)|.$$

Notice, that if the curve X is ordinary and $G = \text{Aut}(X)$ then the cover $X \rightarrow X/G$ is weakly ramified [16, th. 2i] and the above result coincides with the result of G. Cornelissen and F. Kato [5] on deformations of ordinary curves.

4. THE p -RANK REPRESENTATION.

Let D be an effective divisor on a curve X . On the spaces $\Omega_X(-D)$ one can define the action of the Cartier operator. For an introduction to all necessary material the interested reader may consult [20], [25], [13], [24]. There is the following decomposition of the above space

$$\Omega_X(-D) = \Omega_X(-D)^s \bigoplus \Omega_X(-D)^n$$

where $\Omega_X(-D)^s, \Omega_X(-D)^n$ are the spaces of semisimple and nilpotent differentials with respect to the Cartier operator. The above decomposition is compatible with the G -action. For the $k[G]$ -module $\Omega_X(-D)^n$ of nilpotent elements little seems to be known. On the other hand the $k[G]$ -module $V_D := \Omega_X(-D)^s$ was studied by many authors ([13], [7], [3], [24]). The $k[G]$ -module V_D is called in the literature *the p -rank representation*. In general, we have the following decomposition:

$$V_D = \text{core}(V_D) \bigoplus \bigoplus_{S \in \text{Irr} G} P_G(S)^{b(G, D, S)},$$

where S runs over the set of equivalent classes of irreducible representations, $P_G(S)$ denotes the projective cover of S , and $b(G, D, S) \in \mathbb{N}$ are called *the Borne invariants* corresponding to G, D, S [24].

Since G is a p -group and k is assumed to be of characteristic $p > 0$ the only irreducible representation is the trivial one and has projective cover $k[G]$ [21, 15.6]. Moreover $\text{core}(V_{D^*}) = 0$ since D^* is non empty and contains all ramification points [13], [24, 4.5]. Therefore,

$$V_{D^*} = k[G]^{B(G, D^*, k)},$$

where $B(G, D^*, k)$ is an integer.

Proposition 4.1. *With the above notation $B(G, D^*, k) = 2(g_Y - 1) + \gamma_Y - 1 + r$, where g_Y, γ_Y are the genus and the p -rank of the Jacobian of the curve $Y = X/G$, r is the number of points that are ramified in the cover $X \rightarrow Y = X/G$.*

Proof. Let us denote by D_{red}^* the divisor that has the same support with D^* , so that for all prime divisors P we have $v_P(D_{red}^*) > 0 \Rightarrow v_P(D_{red}^*) = 1$. According to [25, p. 175]

$$(18) \quad \dim_k \Omega_X(-D)^s = \dim_k \Omega_X(-D_{red}^*)^s = \dim_k \Omega_X(0)^s + \deg D_{red}^* - 1.$$

But the space $\Omega_X(0)^s$ of semisimple regular differentials is of dimension γ_X , and the degree of D_{red}^* is equal to $2(g_Y - 1)|G| + r$.

The Deuring-Shafarevich formula ([11], [25], [6], [23], [13]) relates the p -ranks γ_X, γ_Y :

$$\gamma_X - 1 = |G| \left(\gamma_Y - 1 + \sum_{i=1}^r \left(1 - \frac{1}{e_0(b_i)} \right) \right).$$

This combined with (18) gives us that

$$\dim_k V_{D^*} = \dim_k \Omega_X(-D^*)^s = |G|(\gamma_Y - 1 + 2(g_Y - 2) + r),$$

and since V_{D^*} is projective we have that $V_{D^*} = k[G]^{\gamma_Y - 1 + 2(g_Y - 2) + r}$, and the desired result follows. $\square$

The above computation allows us to compute the dimension of the space of nilpotent elements. Indeed, the dimension of the space $\Omega_X(-D^*)$ is computed to be equal to

$$\Omega_X(-D^*) = (3g_Y - 3)|G| + |G| \sum_{i=1}^r \sum_{\nu=0}^{\infty} \frac{e_\nu(b_i) - 1}{e_0(b_i)}.$$

Therefore, we have that

$$\dim_k \Omega(-D^*)^n = |G|(g_Y - \gamma_Y) + |G| \sum_{i=1}^r \sum_{\nu=0}^{\infty} \frac{e_\nu(b_i) - 1}{e_0} - r.$$

If the curve X is ordinary *i.e.* $g_Y = \gamma_Y$ then $e_\nu(b_i) = 0$ for all $\nu \geq 2$ and the above formula gives us

$$\dim_k \Omega_X(-D^*)^n = |G| \left(\sum_{i=1}^r \left(1 - \frac{2}{e_0(b_i)} \right) \right).$$

Since this dimension is not divisible by $|G|$ the module $\Omega_X(-D^*)^n$ can not be projective.

The conclusion concerning the dimension is that

$$(19) \quad \dim_k H^1(G, \mathcal{T}_X) = 2(g_Y - 1) + \gamma_Y - 1 + r + \dim_k \Omega_X(-D^*)^n_G.$$

5. BORNE THEORY

Let V_j denote the indecomposable $k[G]$ -module of dimension j . Denote by V the $k[G]$ -module with k -basis $\{e_1, \dots, e_{p^v}\}$ and action given by $\sigma e_\ell = e_\ell + e_{\ell-1}$, $e_0 = 0$. Then, V_j is the subspace of V generated by $\{e_1, \dots, e_j\}$.

Following Borne we define:

Definition 5.1. Let $\pi : X \rightarrow Y$ be a Galois cover of curves defined over k with Galois group $G \cong \mathbb{Z}/p\mathbb{Z}$. For a ramified point P of X we define N_P so that $\sigma(t_P) - t_P$ has valuation $N_P + 1$, where t_P is a local uniformizer at P . Let X_{ram} denote the set of ramification locus of the above cover. For every $0 \leq \alpha \leq p-1$ we define a map $\pi_*^\alpha : \text{Div}_X \rightarrow \text{Div}_Y$ by

$$\pi_*^\alpha D = \left\lfloor \frac{1}{p} \pi_* \left(D - \alpha \sum_{P \in X_{ram}} N_P P \right) \right\rfloor,$$

where $\lfloor \cdot \rfloor$ denotes the integral part of a divisor, taken coefficient by coefficient.

We will use the following

Theorem 5.2. Suppose that X is acted on faithfully by the cyclic p -group $G \cong \mathbb{Z}/p^v\mathbb{Z}$. We break the cyclic extension $X \rightarrow X/G$ to a sequence of cyclic p -extensions by defining for every $1 \leq n \leq v$ the cover $X_n = X/H_n$ where H_n is the unique subgroup of G of order n . We set $X_0 = X$. Let $\pi_n : X_{n-1} \rightarrow X_n$ denote the canonical morphism. Let D be a G -invariant divisor on X , then $H^0(X, \mathcal{O}_X(D)) \cong \bigoplus_{j=1}^v V_j^{\oplus m_j}$. Suppose that $\deg(D) \geq 2g_X - 2$. Then, the integers m_j are given by

$$m_j = \deg(\pi_{v*}^{\alpha_0(j)} \dots \pi_{1*}^{\alpha_{v-1}(j)} D) - \deg(\pi_{v*}^{\alpha_0(j+1)} \dots \pi_{1*}^{\alpha_{v-1}(j+1)} D) \text{ if } 1 \leq j \leq p^v - 1,$$

$$m_{p^v} = 1 - g_{X_v} + \deg(\pi_{v*}^{p-1} \dots \pi_{1*}^{p-1} D),$$

where for $1 \leq j \leq p^v$ the integers $\alpha_0(j), \dots, \alpha_{v-1}(j)$ are the digits of the p -adic expansion of $j-1$ i.e.,

$$j-1 = \sum_{h=0}^{v-1} \alpha_h(j) p^h, \text{ with } 0 \leq \alpha_h(j) \leq p-1.$$

Proof. [4, th. 7.25] □

We would like to consider the space $\Omega_X(-D^*) = \mathcal{O}_X(K + D^*) = \mathcal{O}_X(2D^*)$, where $D^* = \pi^*(\text{div}(\phi)) + \sum_{P \in X_{ram}} \sum_{i=0}^{\infty} (e_i(P) - 1)P$. Notice that $\deg(2D^*) = 4g_X - 4 \geq 2g_X - 2$ so theorem 5.2 is applicable.

Define

$$D(j) := \pi_{v*}^{\alpha_0(j)} \dots \pi_{1*}^{\alpha_{v-1}(j)} 2D^*.$$

We have $m_j = \deg(D(j)) - \deg(D(j+1))$ for $1 \leq j \leq p^v - 1$. Since $\dim_k H_0(G, V_j) = \dim_k (V_j)_G = 1$ the desired dimension is given by:

$$(20) \quad \dim_k \mathcal{O}_X(2D^*)_G = \sum_{j=1}^{p^v} m_j = \deg(D(1)) - \deg(D(p^v)) + 1 - g_X + \deg(\pi_{v*}^{p-1} \dots \pi_{1*}^{p-1} 2D^*).$$

Observe that the p -adic expansion of $p^v - 1$ is

$$p^v - 1 = p - 1 + (p-1)p + \dots + (p-1)p^{v-1},$$

so $\alpha_i(0) = 0, \alpha_i(p^v - 1) = p - 1$ for all $0 \leq i \leq v - 1$. Therefore (20) is simplified to

$$(21) \quad \dim_k \mathcal{O}_X(2D^*)_G = \deg(D(1)) + 1 - g_X.$$

Definition 5.3. Let $X_{ram,i}$ be the set of points of X_i ramified in the cover $X_i \rightarrow X_{i+1}$.

Notice that $\mathbb{Z}/p\mathbb{Z} = \text{Gal}(X_0/X_1)$ is contained in every subgroup of G , therefore $X_{ram} = X_{ram,0}$.

Denote by Δ the map $\mathbb{Z} \rightarrow \mathbb{Z}$ sending $\Delta : a \mapsto \left\lfloor \frac{1}{p}a \right\rfloor$, and by Δ^n the composition $\Delta \circ \dots \circ \Delta$ of n times Δ . If $a = \sum_{i=0}^{\ell} \alpha_i p^i$ is the p -adic expansion of a , then

$$\Delta^n(a) = \sum_{i=n}^{\ell} \alpha_i p^{i-n}.$$

Assume that $H = \mathbb{Z}/p\mathbb{Z}$. Let $O(P) = \{gP : g \in H \setminus H(P)\}$ be the orbit of P under the action of H , and consider the H -invariant divisor: $\sum_{Q \in O(P)} aQ$. We have that

$$\pi_*^1 \left(\sum_{Q \in O(P)} aQ \right) = \begin{cases} a \cdot \pi_*(Q) & \text{if } Q \text{ does not ramify in } X \mapsto X/G \\ \Delta(a) \cdot \pi_*(Q) & \text{if } Q \text{ ramifies in } X \mapsto X/G \end{cases}$$

The divisor $2D$ can be written as

$$2\pi^*\text{div}(\phi) + \sum_{P \in X_{ram}} \sum_{i=0}^{\infty} (2e_i(P) - 2)P.$$

Since the divisor $\pi^*\text{div}(\phi)$ has empty intersection with the ramification locus, we have

$$\begin{aligned} \pi_*^1(2D^*) &= (\pi_v \circ \pi_{v-1} \circ \dots \circ \pi_2)^* 2\text{div}(\phi) + \sum_{P \in \pi_{1*}(X_{ram})} \left[\frac{1}{p} \sum_{i=0}^{N_P} (2e_i(P) - 2) \right] P = \\ &= (\pi_v \circ \pi_{v-1} \circ \dots \circ \pi_2)^* 2\text{div}(\phi) + \sum_{P \in \pi_{1*}(X_{ram})} \Delta \left(\sum_{i=0}^{N_P} (2e_i(P) - 2) \right) P. \end{aligned}$$

The decomposition group $G_0(P)$ is a cyclic subgroup of the whole group G , therefore $G_0(P) = \mathbb{Z}/p^{k(P)}\mathbb{Z}$. Observe that a point $P \in X_{ram}$ is fully ramified in all covers $X_i \rightarrow X_{i+1}$ for $i \leq k(P)$. We can see that

$$D(1) = 2\text{div}(\phi) + \sum_{P \in \pi_*(X_{ram})} \sum_{i=1}^{k(P)} \Delta^{k(P)} \left(\sum_{i=0}^{N_P} (2e_i(P) - 2) \right) P.$$

Let

$$1 \leq i_1 \leq i_2 \leq \dots \leq i_{k(P)} = N_P,$$

be the sequence of the jumps in the ramification filtration at the point P , i.e.

$$G_{i_k} \supsetneq G_{i_k+1} \text{ and } G_{i_{k(P)}+1} = \{1\}.$$

Lemma 5.4. There are strictly positive integers $a_0, a_1, \dots, a_{k-1}$, so that the sequence of jumps for the ramification filtration for the cyclic group $\mathbb{Z}/p^k\mathbb{Z}$ is of the form:

$$i_\nu = \sum_{\mu=0}^{\nu-1} a_\mu p^\mu.$$

In particular, for all $\nu \geq 2$

$$i_\nu - i_{\nu-1} = a_{\nu-1}p^{\nu-1}$$

Proof. This is a direct consequence of the Hasse-Arf theorem for Abelian groups and it is explained in the example that appears on page 76 in [22]. $\square$

Notice that we have $k(P)$ jumps in the ramification filtration since G_{i_k}/G_{i_k+1} is elementary Abelian, therefore isomorphic to $\mathbb{Z}/p\mathbb{Z}$. We set $d_P = \sum_{i=0}^{N_P} (2e_i(P) - 2)$, and we compute:

$$d_P = \sum_{i=0}^{i_1} (2e_{i_1}(P) - 2) + \sum_{i=i_1+1}^{i_2} (2e_{i_2}(P) - 2) + \cdots + \sum_{i_{k(P)-1}+1}^{i_{k(P)}} (2e_{i_{k(P)}}(P) - 2).$$

Since $e_{i_\ell} = p^{k(P)-\ell+1}$ for all $1 \leq \ell \leq k(P)$ we have that

$$\begin{aligned} d_P &= 2(p^{k(P)} - 1)(i_1 + 1) + 2((p^{k(P)-1} - 1)(i_2 - i_1) + \cdots + 2(p-1)(i_{k(P)} - i_{k(P)-1})) = \\ &= \sum_{\nu=1}^{k(P)} 2(p^{k(P)-\nu+1} - 1)(i_\nu - i_{\nu-1}), \text{ where } i_0 = -1. \end{aligned}$$

Using lemma 5.4 we obtain that there are integers $a_0(P), a_1(P), \dots, a_{k(P)-1}(P)$ so that

$$\begin{aligned} (22) \quad d_P &= 2p^{k(P)} - 2 + \sum_{\nu=1}^{k(P)} 2p^{k(P)-\nu+1}p^{\nu-1}a_{\nu-1}(P) + \sum_{\nu=1}^{k(P)} 2(-i_\nu + i_{\nu-1}) = \\ &= 2p^{k(P)} + p^{k(P)}2 \sum_{\nu=1}^{k(P)} a_{\nu-1} - 2N_P - 2 \end{aligned}$$

One can compute [22, exam. p.72] that the sum $f(P) := \sum_{\nu=1}^{k(P)} a_{\nu-1}(P)$ is the highest jump in the upper ramification filtration. Using (22) we compute that

$$\Delta^{k(P)}(d_P) = 2f(P) + \Delta^k(-2N_P - 2).$$

Combining all the above we arrive at

$$(23) \quad \deg D(1) = 4g_Y - 4 + 2\#\pi_*(X_{ram}) + \sum_{P \in \pi_*(X_{ram})} (2f(P) + \Delta^k(-2N_P - 2))$$

The desired result follows:

$$\dim_k \mathcal{O}(2D^*) = 3(g_Y - 1) + 2\#\pi_*(X_{ram}) + \sum_{P \in \pi_*(X_{ram})} (2f(P) + \Delta^k(-2N_P - 2)).$$

Observe that if $G = \mathbb{Z}/p\mathbb{Z}$ the above result coincides with the computation of [1],[10].

Example: Let $\sigma_i = \sum_{\nu=0}^{i-1} a_\nu$ $i = 0, \dots, k(P)$ be the jumps in the upper ramification filtration at the wild ramified point P . It is known that $\sigma_{i+1} = p\sigma_i$ or $\sigma_{i+1}/p \geq \sigma_i$ and $p \nmid \sigma_{i+1}$ [18],[19]. Assume that all jumps in the ramification filtration are of the form $\sigma_{i+1} = p\sigma_i$. In this case one can prove by induction that $a_i = a_0p^{i-1}(p-1)$, and we can compute

$$f(P) + \Delta^k(-2N_P - 2) = \sum_{\nu=0}^{k(P)-1} a_\nu + \left[\frac{-2 \sum_{\nu=0}^{k(P)-1} a_\nu p^\nu - 2}{p^{k(P)}} \right] =$$

$$\begin{aligned}
&= \sum_{\nu=0}^{k(P)-1} a_0(p-1)p^{\nu-1} + \left\lfloor \frac{-2 \sum_{\nu=0}^{k(P)-1} (p-1)p^{2\nu-1} - 2}{p^{k(P)}} \right\rfloor \\
&= a_0(p^{k(P)} - 1) + \left\lfloor -a_0 2 \frac{p^{2k(P)} - 1}{p^{k(P)+1}(p+1)} - \frac{2}{p^{k(P)}} \right\rfloor.
\end{aligned}$$

REFERENCES

1. José Bertin and Ariane Mézard, *Déformations formelles des revêtements sauvagement ramifiés de courbes algébriques*, Invent. Math. **141** (2000), no. 1, 195–238. MR 2001f:14023
2. Niels Borne, *Une formule de riemann-roch équivariante pour les courbes*, Ph.D. thesis, Laboratoire de Mathématiques Pures de l' Université Bordeaux, 2000.
3. ———, *A relative Shafarevich theorem*, Math. Z. **248** (2004), no. 2, 351–367. MR MR2088933 (2005g:14058)
4. ———, *Cohomology of G -sheaves in positive characteristic*, Adv. Math. **201** (2006), no. 2, 454–515. MR MR2211535
5. Gunther Cornelissen and Fumiharu Kato, *Equivariant deformation of Mumford curves and of ordinary curves in positive characteristic*, Duke Math. J. **116** (2003), no. 3, 431–470. MR MR1958094 (2004c:14044)
6. Max Deuring, *Automorphismen und Divisorenklassen der Ordnung ι in algebraischen Funktionenkörpern*, Math. Ann. **113** (1937), no. 1, 208–215. MR MR1513087
7. Ernst Kani, *The Galois-module structure of the space of holomorphic differentials of a curve*, J. Reine Angew. Math. **367** (1986), 187–206. MR 88f:14024
8. Bernhard Köck, *Galois structure of Zariski cohomology for weakly ramified covers of curves*, Amer. J. Math. **126** (2004), no. 5, 1085–1107. MR MR2089083 (2005i:11163)
9. Aristides I. Kontogeorgis, *On the tangent space of the deformation functor of curves with automorphisms*, arXiv:math.AG/0201194 v3 2005.
10. ———, *Polydifferentials and the deformation functor of curves with automorphisms*, to appear in the JPAA.
11. Manohar L. Madan, *On a theorem of M. Deuring and I. R. Šafarevič*, Manuscripta Math. **23** (1977/78), no. 1, 91–102. MR MR0460335 (57 #329)
12. Shōichi Nakajima, *On Galois module structure of the cohomology groups of an algebraic variety*, Invent. Math. **75** (1984), no. 1, 1–8. MR 85j:14021
13. ———, *Equivariant form of the Deuring-Šafarevič formula for Hasse-Witt invariants*, Math. Z. **190** (1985), no. 4, 559–566. MR 87g:14024
14. ———, *Action of an automorphism of order p on cohomology groups of an algebraic curve*, J. Pure Appl. Algebra **42** (1986), no. 1, 85–94. MR MR852320 (88d:14018)
15. ———, *Galois module structure of cohomology groups for tamely ramified coverings of algebraic varieties*, J. Number Theory **22** (1986), no. 1, 115–123. MR 87i:14010
16. ———, *p -ranks and automorphism groups of algebraic curves*, Trans. Amer. Math. Soc. **303** (1987), no. 2, 595–607. MR 88h:14037
17. Rachel J. Pries, *Equiramified deformations of covers in positive characteristic*, arXiv:math.AG/0403056 v3.
18. ———, *Wildly ramified covers with large genus*, J. Number Theory **119** (2006), no. 2, 194–209. MR MR2250044
19. Hermann Ludwig Schmid, *Zur Arithmetik der zyklischen p -Körper.*, J. Reine Angew. Math. **176** (1936), 161–167 (German).
20. Jean-Pierre Serre, *Sur la topologie des variétés algébriques en caractéristique p* , Symposium internacional de topología algebraica International symposium on algebraic topology, Universidad Nacional Autónoma de México and UNESCO, Mexico City, 1958, pp. 24–53. MR MR0098097 (20 #4559)
21. ———, *Linear representations of finite groups*, Springer-Verlag, New York, 1977, Translated from the second French edition by Leonard L. Scott, Graduate Texts in Mathematics, Vol. 42. MR MR0450380 (56 #8675)
22. ———, *Local fields*, Springer-Verlag, New York, 1979, Translated from the French by Marvin Jay Greenberg. MR 82e:12016
23. I. Shafarevitch, *On p -extensions*, Rec. Math. [Mat. Sbornik] N.S., AMS Transl. series II **4** (1954), 59–72. MR MR0020546 (8,560e)

24. Nicolas Stalder, *On p -rank representations*, J. Algebra **280** (2004), no. 2, 825–841. MR MR2090066 (2005h:14074)
25. Doré Subrao, *The p -rank of Artin-Schreier curves*, Manuscripta Math. **16** (1975), no. 2, 169–193. MR MR0376693 (51 #12868)
26. Charles A. Weibel, *An introduction to homological algebra*, Cambridge University Press, Cambridge, 1994. MR 95f:18001

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF THE ÆGEAN, 83200 KARLOVASSI, SAMOS, GREECE, <http://eloris.samos.aegean.gr>
E-mail address: kontogar@aegean.gr